



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Full-length nucleotide sequences of mcr-1 harboring plasmids isolated from extended-spectrum β -lactamase (ESBL)- producing *Escherichia coli* of different origins

Zurfluh, Katrin ; Klumpp, Jochen ; Nüesch-Inderbinen, Magdalena ; Stephan, Roger

Abstract: Here, we present the full sequences of three mcr-1-carrying plasmids isolated from extended-spectrum- β -lactamase (ESBL)-producing *Escherichia coli*. The plasmids belong to three different replicon types and are 34,640 bp, 209,401 bp, and 247,885 bp in size. We describe for the first time a composite transposon containing mcr-1 localized on a multidrug-resistant (MDR) IncHI2 plasmid harboring additional determinants of resistance to six different classes of antibiotics, including the ESBL gene blaCTX-M-1, and heavy metal resistance.

DOI: <https://doi.org/10.1128/AAC.00935-16>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-129982>

Journal Article

Accepted Version

Originally published at:

Zurfluh, Katrin; Klumpp, Jochen; Nüesch-Inderbinen, Magdalena; Stephan, Roger (2016). Full-length nucleotide sequences of mcr-1 harboring plasmids isolated from extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* of different origins. *Antimicrobial Agents and Chemotherapy*, 60(9):5589-5591. DOI: <https://doi.org/10.1128/AAC.00935-16>

Full-length nucleotide sequences of *mcr-1* harboring plasmids isolated from extended-spectrum β -lactamase (ESBL)- producing *Escherichia coli* of different origins

Katrin Zurfluh^{1#}, Jochen Klumpp², Magdalena Nüesch-Inderbinen¹, Roger Stephan¹

¹Institute for Food Safety and Hygiene, Vetsuisse Faculty University of Zurich, Winterthurerstr. 272, CH-8057 Zurich, Switzerland

²Institute of Food, Nutrition and Health, ETH Zurich, Schmelzbergstr. 7, CH-8092 Zurich, Switzerland

Running Title: Nucleotide sequences of *mcr-1* harboring plasmids

Corresponding author:

Katrin Zurfluh, Institute for Food Safety and Hygiene, Vetsuisse Faculty University of Zurich, Winterthurerstr. 272, CH-8057 Zurich, Switzerland, Phone 0041-44-6358651, Fax 0041-44-6358908, e-mail katrin.zurfluh@uzh.ch

Abstract

Here we present the full sequences of three *mcr-1* carrying plasmids isolated from ESBL-producing *E. coli*. The plasmids belong to three different replicon types and are 34.640 bp, 209.401 bp and 247.885 bp in size, respectively. We describe for the first time a composite transposon containing *mcr-1* localized on an MDR-IncHI2 plasmid harboring additional determinants of resistance to six different classes of antibiotics including the ESBL-gene *bla*_{CTX-M-1} and heavy metal resistance.

The recent description of the plasmid-mediated colistin resistance gene, *mcr-I*, in strains isolated from food animals, food and humans in China was a trigger for an avalanche of retrospective studies investigating the presence of this specific gene (1). Since then, *mcr-I* has been found almost all over the world and the earliest evidence for its presence dates back to the 1980's (2). The wide spread of *mcr-I* and the finding that this resistance marker is often associated with multidrug resistant *Enterobacteriaceae*, e.g. extended-spectrum β -lactamase (ESBL)-producers or carbapenemase-producers, is of great concern to public health (3). The *mcr-I* gene was so far associated with different plasmid replicon types such as IncI2, IncHI2, IncP, IncFIP and IncX4 (1,4,5,6,7). However, only very limited data about the complete sequences of such plasmids and the genetic structure surrounding the *mcr-I* gene are available. In this study, we present the complete sequence of three distinct *mcr-I* harboring plasmids isolated from ESBL-producing *E. coli* isolates originating from river water (*E. coli* OW3E1, B1:ST359; SHV-12), imported vegetables from Thailand (*E. coli* H226B, A:ST167, CTX-M-55) and imported poultry meat from Italy (*E. coli* S38, B1:ST602, CTX-M-1) (8, 9).

The *mcr-I* harboring plasmids were transferred by transformation experiments into *E. coli* DH5- α and colistin resistant transformants were selected on LB-agar supplemented with 2 mg/L colistin (SIGMA, St. Louis, USA). The *mcr-I* harboring plasmids pOW3E1, pH226B and pS38 were extracted using the Large-Construct Kit (Qiagen, Hombrechtikon, Switzerland) according the manufacturer's protocol. The plasmids were sequenced on a PacBio RS2 device (Pacific Biosciences, Menlo Park, USA) with a 10 kb size-selected insert library and P6/C4 chemistry. pOW3E1 and pH226B were multiplexed with four other plasmids (data not shown) in a barcoded library (symmetric 384 barcode set) and run on two SMRT cells; pS38 was prepared as single library and sequenced on another SMRT cell.

De-Barcoding, whitelisting and *de novo* assembly (HGAP3 algorithm) was performed using SMRTanalysis version 2.3 (Pacific Biosciences). HGAP3 settings were kept at the defaults, except for the expected genome size, which was set between 60 and 200. The HGAP3 analysis produced complete plasmid sequences of the multiplexed samples with at least 50-fold coverage over the entire molecules. Plasmid pS38 was sequenced individually with 2250-fold coverage. The plasmid sequence was automatically annotated using the online Rapid Annotation Subsequencing Technology (RAST) and CLC Main Workbench Version 7.7 (CLC bio, Aarhus, Denmark) (10). Automated annotation was manually refined using the BLASTn and BLASTp programs (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Plasmid pOW3E1 is 34,640 bp in size, with a G+C content of 42.3% (Table 1). The plasmid backbone is similar to that of other IncX4 plasmids including a type IV secretion system (*pilX* operon), which is associated with this plasmid group (data not shown). Interestingly, no resistance determinants other than the *mcr-I* gene were located on pOW3E1.

Plasmid pH226B is 209,401 bp in size with a G+C content of 46.5% (Table 1). It consists of a typical IncHI1 backbone and shares many open reading frames (orf) with the reference plasmid of the IncHI1 group R27 (GenBank accession number AF250878). Beside the *mcr-I* gene, a Tn3 family transposon carrying heavy metal resistance determinants (e.g. Zn^{2+} , Cu^{2+} or Ag^{2+}) is present on pH226B as well as a remotely located *tcuABC* operon that enables the use of tricarballylate as carbon/energy source.

Plasmid pS38, extracted from an *E. coli* isolate originating from imported poultry meat, represents a large multi-drug-resistant (MDR) plasmid (247,885 bp, G+C content: 46.7%) with high similarity in its backbone to the IncHI2 reference plasmid R478 (accession no. NC_005211) (Table 1). The MDR-region of this plasmid includes two class 1 integrons carrying transposons In641 (*estX-3*, *psp*, *aadA2*, *cmlA1*, *aadA1a*, *qacH2*) and In369 (*dfrA1b*, *aadA1b*), conferring resistance to aminoglycosides, chloramphenicol, quaternary ammonium compounds and

76 trimethoprim. Between these transposons, the macrolide efflux pump gene *mefB*, the ESBL-gene
77 *bla*_{CTX-M-1} and the sulphonamide resistance gene *sul3* are present. Downstream of transposon
78 In369, two transposons carrying mercury resistance determinants and tetracycline resistance
79 determinants, respectively, followed. Furthermore, a tellurite resistance operon is present on
80 pS38.

81 The *mcr-1* genes were found to be located within various genetic contexts in the sequenced
82 plasmids (Figure 1). In all three plasmids, a 735 bp orf encoding a hypothetical protein with
83 similarities to a PAP2 superfamily protein was detected immediately downstream of the *mcr-1*
84 gene (both together hereafter referred to as *mcr-1* element).

85 In pOW3E1 the *mcr-1* gene is inserted approximately 1.5 kb upstream of the replicon initiation
86 protein and is found close to an IS2-like insertion element. An IS*AplI* element is absent. In
87 pH226B, the *mcr-1* element is embedded in the IncHI1 backbone between two genes that were
88 highly similar to those annotated as Orf061 and Orf062 on the IncHI1 reference plasmid R27.
89 No mobile element was present upstream or downstream nearby. In pS38, the *mcr-1* element is
90 flanked by two IS*AplI* elements (composite transposon) and is located directly upstream of the
91 MDR-region.

92 To our knowledge, this is the first description of an MDR-IncHI2 plasmid harboring a composite
93 transposon containing the *mcr-1* gene and determinants for resistance to six different classes of
94 antibiotics including the ESBL-gene *bla*_{CTX-M-1} and additional heavy metal resistance
95 determinants. The fact that the *mcr-1* gene is found to be part of a composite transposon is
96 highly worrisome, since accelerated dissemination by illegitimate recombination of the whole
97 transposon to either plasmids or chromosomes is likely.

98 Furthermore, we present for the first time an IncHI1 plasmid carrying the mobile colistin
99 resistance determinant. The presence of heavy metal resistance determinants on pS38 and

pH226B might lead to co-selection, for example by zinc which is widely used as a feed additive for fattening pigs.

Since the *mcr-I* element was found on different plasmid types and the IS*AplI* element was not always present we hypothesize that the *mcr-I* gene may have been mobilized independently several times. In the case of pH226B however, where the *mcr-I* gene is embedded in the middle of the backbone without any evidence for the presence of mobile elements, no conclusions concerning the mobilization of this gene can be made.

Nucleotide sequence

The GenBank accession numbers for pS38, pOW3E1 and pH226B are KX129782, KX129783, KX129784, respectively.

Acknowledgments

We are grateful to the staff of the Functional Genomics Center Zurich, particularly Anna Bratus, Andrea Patrignani and Weihong Qi for excellent technical support and bioinformatics support.

References

1. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2015. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: A microbiological and molecular biological study. Lancet Infect. Dis. **16**: 161–168. [http://dx.doi.org/10.1016/S1473-3099\(15\)00424-7](http://dx.doi.org/10.1016/S1473-3099(15)00424-7).
2. Shen Z, Wang Y, Shen Y, Shen J, Wu C. 2016. Early emergence of *mcr-I* in *Escherichia coli* from food-producing animals. Lancet Infect Dis. **16**:293. [doi: 10.1016/S1473-3099\(16\)00061-X](https://doi.org/10.1016/S1473-3099(16)00061-X)

3. **Falgenhauer L, Waezsada SE, Yao Y, Imirzalioglu C, Käsbohrer A, Roesler U, Michael GB, Schwarz S, Werner G, Kreienbock L, Chakraborty T, for the RESET consortium.** 2016. Colistin resistance gene *mcr-I* in extended-spectrum beta-lactamase-producing and carbapenemase-producing Gram-negative bacteria in Germany. *Lancet Infect Dis.* **16**:282-3. doi: 10.1016/S1473-3099(16)00009-8.
4. **Zhi C, Lv L, Yu LF, Doi Y, Liu JH.** 2016. Dissemination of the *mcr-I* colistin resistance gene. *Lancet Infect Dis.* **16**:292-293. doi: 10.1016/S1473-3099(16)00063-3.
5. **Nordmann P, Lienhard R, Kieffer N, Clerc O, Poirel L.** 2016. Plasmid-mediated colistin-resistant *Escherichia coli* in bacteremia in Switzerland. *Clin Infect Dis.* **15**:1322-3. doi: 10.1093/cid/ciw124.
6. **Doumith M, Godbole G, Ashton P, Larkin L, Dallman T, Day M, Day M, Muller-Pebody B, Ellington MJ, de Pinna E, Johnson AP, Hopkins KL, Woodford N.** 2016. Detection of the plasmid-mediated *mcr-I* gene conferring colistin resistance in human and food isolates of *Salmonella enterica* and *Escherichia coli* in England and Wales. *J. Antimicrob. Chemother.*, accepted pii: dkw093.
7. **Li A, Yang Y, Miao M, Chavda KD, Mediavilla JR, Xie X, Feng P, Tang YW, Kreiswirth BN, Chen L, Du H.** 2016. Complete sequences of *mcr-I* -harboring plasmids from extended spectrum β -lactamase (ESBL)- and carbapenemase-producing Enterobacteriaceae (CPE). *Antimicrob. Agents Chemother.* accepted. AAC.00550-16. <http://aac.asm.org/lookup/doi/10.1128/AAC.00550-16>.
8. **Zurfluh K, Poirel L, Nordmann P, Nüesch-Inderbinen M, Hächler H, Stephan R.** 2016. Occurrence of the plasmid-borne *mcr-I* colistin resistance gene in extended-spectrum- β -lactamase-producing Enterobacteriaceae in river water and imported vegetable samples in Switzerland. *Antimicrob. Agents Chemother.* **60**: 2594-2595. <http://aac.asm.org/lookup/doi/10.1128/AAC.00066-16>.

9. **Zogg AL, Zurfluh K, Nüesch-Inderbinen M, Stephan R.** 2016. Characteristics of
ESBL-producing Enterobacteriaceae and Meticillin-resistant *Staphylococcus aureus*
(MRSA) isolated from Swiss and imported raw poultry meat collected at retail level.
Schweiz. Arch. Tierheilk., in press.
10. **Aziz RK, Bartles D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes
S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA,
McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R,
Vassieva O, Vonstein V, Wilke A, Zagnitko O.** 2008. The RAST Server: rapid
annotations using subsystems technology. BMC Genomics. 9:75. doi:10.1186/1471-2164-
9-75.

Fig 1. The genetic structures surrounding the *mcr-I* gene in plasmids of this study compared to
plasmid pHNSHP45. The arrows indicate open reading frames (orf), with black, stippled, grey
and white arrows representing *mcr-I*, IS elements, orf with known function and orf with
unknown function, respectively.